

Anticipatory VtoV coarticulation in French in several Motor Speech Disorders

Daria D'Alessandro¹, Michaela Pernon^{1,2}, Cécile Fougeron¹ and Marina Laganaro³

¹Laboratoire de Phonétique et Phonologie, UMR 7018, CNRS/U. Sorbonne-Nouvelle

²Dpt des Neurosciences Cliniques, HUG

³FAPSE, Université de Genève

V-to-V anticipatory coarticulation is frequently taken as an indicator of the anticipated planning of an up-coming vowel, and thus the manifestation of a process where context-dependent speech targets or coordinated gestures are planned together before execution (e.g. [1], [2], [3]). In this study, V-to-V anticipatory coarticulation in French is compared in 4 groups of patients presenting different Motor Speech Disorders --Apraxia of Speech (AoS) and 3 types of dysarthrias associated with Amyotrophic Lateral Sclerosis (D-ALS), Parkinson (D-Pk) and Wilson (D-WI) diseases-- with the aim of getting further insights into the understanding of the control of coarticulation in speech (as in [4], [5], [6]). It is hypothesized that perturbed coarticulatory patterns may arise as a consequence of disruption in the planning and sequencing of speech targets into cohesive speech units, as expected in AoS where speech is often described as 'syllabified' or 'segmentalized'. However, it can also derive from deficits in the motor programming or execution of speech whose consequences are slower movements, restricted displacement and/or perturbed coordination between gestures, as can be encountered in dysarthria. So far, inconsistent results have been found in the literature regarding coarticulatory patterns in different MSD (see [5] & [6]). Moreover, AoS patients are more often compared to patients with aphasia than to patients with dysarthria. More data is thus needed to understand whether deficits linked to distinct cerebral disorders do impact coarticulation, and how they can shed light on the levels and units over which coarticulation is planned.


Twenty patients (five per groups) suffering from mild to moderate speech impairments, and 40 healthy control speakers (age and sex matched) were recorded while reading a short text containing twelve disyllabic [pV₁pV₂] French words with V₁=/a/ and V₂=/a/ or /i/. The pivot bilabial consonant /p/ was chosen in order to maximize the effect of V₂ on V₁ shown by a lowering of F1 and a rising of F2, leading to a less compact F2-F1, when /a/ is followed by /i/.

Results show a significant contextual effect of V₂ on V₁ on F2-F1 in all groups of patients with dysarthria and in the control group, but not in the group of patients with AoS (Fig. 1), as found in [8], for instance. Nonetheless, some contextual effect is also present for some speakers in the AoS group, with a certain speaker- and token-dependent variability (Fig.3). In order to compare further the groups, the amount of coarticulation in the /papi/ (V₂= /i/) words was computed as the ratio between the F2 of the first vowel /a/ and the F2 of the following /i/. Reduced amount of coarticulation (more distance between [a] and [i] F2's) is found in both the D-WI group and AoS groups, compared to the CTRL and other MSD groups (Fig. 2).

Altered coarticulatory patterns (absent or reduced) for the patients in the AoS and D-WI groups can both be related to deficits in speech coordination and timing, but over different domains. In the Wilson group, reduced coarticulation is associated with longer V₁ and with restricted articulatory movements' displacement, as shown by a reduced acoustic space for all Wilson patients especially in the F1 dimension (linked to reduced jaw displacements). Taken together this could suggest that it is at the segment level that spatio-temporal coordination is altered for these speakers, with a pressure towards articulatory accuracy rather than speed, and potentially less overlap between V₁ and V₂. The AoS group presents longer V₁ targets but also longer intervals between V₁ and V₂, due to either long pivot Cs and/or internal pauses, suggesting an altered phasing between the two syllables in the [pV₁.pV₂] word. Together with the lack of anticipation of the up-coming V₂ during the first syllable, AoS

patients seem to proceed with syllable-sized phonetic plans lacking inter-syllable coordination.

While more speakers per patient groups are needed, and local anticipatory coarticulation remains to be examined, theoretical implications of these results will be discussed at the conference.



Apraxia of Speech (AoS) ; Dysarthria associated with Wilson disease (D-WI), with Amyotrophic Lateral Sclerosis ((D-ALS); with Parkinson disease (D-Pk) ; healthy control speakers (CTRL)

Figure 1. Acoustic compacity of $V_1=/a/$, measured as F_2-F_1 , per context ($V_2=/a/$ or $V_2=/i/$) and per group




Figure 2. Degree of coarticulation, measured by the ratio of F_2 in $V_1/a/$ and $V_2/i/$ in $/api/$ words, per group (smaller ratio = less coarticulation)




Figure 3. F_1 and F_2 of $/a/$ according to V_2 context ($/a/$ or $/i/$) for the 5 patients with Apraxia of Speech (different color per patient, 12 tokens each, F=female, H= male speakers).

References:

- [1] Whalen, D. H. 1990. Coarticulation is largely planned. *Journal of Phonetics*, 18, 3-35.
- [2] Bell-Berti, F., Krakow, R. A., Gelfer, C. E., & Boyce, S. E. 1995. Anticipatory and carryover effects: Implications for models of speech production. In F. Bell-Berti & L.J. Raphael (Eds.), *Producing speech: contemporary issues for Katherine Safford Harris*, 77-98.
- [3] Recasens, D. 2002. An EMA study of VCV coarticulatory direction. *The Journal of the Acoustical Society of America*, 111(6), 2828-2841.
- [4] Hertrich, I., & Ackermann, H. 1999. Temporal and spectral aspects of coarticulation in ataxic dysarthria: An acoustic analysis. *Journal of Speech, Language, and Hearing Research*, 42(2), 367-381.
- [5] Katz, W.F. 2000. Anticipatory coarticulation and aphasia: implications for phonetic theories. *Journal of Phonetics*, 28, 313-334

- [6] Frisch, S. A., Maxfield, N., & Belmont, A. 2016. Anticipatory coarticulation and stability of speech in typically fluent speakers and people who stutter. *Clinical linguistics & phonetics*, 30(3-5), 277-291.
- [7] Hardcastle, B., & Tjaden, K. 2008. Coarticulation and Speech Impairment. In. M. J. Ball, M.R. Perkins, N. Müller, S. Howard, eds., *The handbook of clinical linguistics*. Blackwell.
- [8] Ziegler, W., & von Cramon, D. (1986). Disturbed coarticulation in apraxia of speech : acoustic evidence. *Brain and Language*, 29, 34-47.